

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Numéro de publication

0 000 322  
A1

(12)

## DEMANDE DE BREVET EUROPEEN

(21) Application number: 78850006.4

(22) Date of filing: 03.07.78

(51) Int. Cl.<sup>2</sup>: C 07 C 87/28, C 07 C 87/29,  
C 07 C 93/14, C 07 C 49/16,  
C 07 C 131/00, C 07 C 49/06,  
C 07 C 91/18, // C 07 C 125/00,  
C 07 C 25/12

(30) Priority: 04.07.77 GB 27992/77  
31.05.78 GB 27992/77  
22.05.78 GB 21249/78

(43) Date of publication of application:  
10.01.79 Bulletin 79/1

(84) Designated Contracting States:  
BE CH DE FR LU NL SE

(71) Applicant: Astra Läkemedel Aktiebolag,  
S-151 85 Södertälje (SE)

(72) Inventor: Carnmalm, Bernt Sigfrid Emanuel,  
Törnrosavägen 14,  
S-151 52 Södertälje (SE)

(72) Inventor: Högberg, Thomas,  
Vallmostigen 5,  
S-150 20 Järna (SE)


(72) Inventor: de Paulis, Tomas,  
Villavägen 13,  
S-150 11 Björnlunda (SE)

(72) Inventor: Ross, Svante Bertil,  
Hedvägen 8,  
S-151 52 Södertälje (SE)

(72) Representative: Wurm, Bengt R. et al,  
S-151 85 Södertälje (SE)

(54) Compounds having an anti-depressive or tranquilizing activity, pharmaceutical compositions containing them, and processes and intermediates for their preparation.

(57) Compounds of the formula



wherein the dotted line represents an optional double bond  
and Ar represents the group



wherein Y is bound in the 2-, 3- or 4-position and represents a lower alkyl or lower alkoxy group, a halogen, a trifluoromethyl group or an amino, mono- or di-lower alkylamino group, or Ar represents a pyridyl group bound in the 2-, 3- or 4-position, X represents hydrogen, a lower alkyl or lower alkoxy group, a halogen, a trifluoromethyl group or an amino, mono- or di-lower alkylamino group, R represents a lower alkyl group, and R' represents hydrogen or a lower alkyl group, or a therapeutically acceptable acid addition salt thereof, or a bioprecursor thereof.

The compounds are prepared by various methods; pharmaceutical compositions containing the compounds are described.

0000322

Astra Läkemedel Aktiebolag  
Södertälje/SWEDEN

Inventors: B S E Carnmalm, T Högberg, T de Paulis, S B Ross

LA 534-1

78 06 19

IH/ACE

Compounds Having Anti-Depressive or Tranquilizing Activity,  
Pharmaceutical Compositions Containing Them, and Processes  
and Intermediates for Their Preparation

The present invention is related to new compounds of the  
diarylalkylamine type having therapeutic activity, to methods  
5 for preparing such compounds, to pharmaceutical preparations  
comprising such compounds and to methods of treatment  
employing such compounds. The invention is also related to  
intermediates useful in preparation of such therapeutically  
active compounds.

10

The object of the invention is to obtain compounds having a  
therapeutical activity in the central nervous system, especi-  
ally an anti-depressive or a tranquilizing activity.

15 Prior Art

British Patent 1 429 068 discloses compounds corresponding to  
the general formula:



0000322.



10 closes compounds of the general formula:



20 having anti-depressive activity.

Compounds following within the general formula



35 =OH,  $R^{III}=R^{IV}=H$ , and  $R^I=Cl$ ,  $R^{II}=OH$ ,  $R^{III}=CH_3$ ,  $R^{IV}=H$  having hypoc-

tensive properties, by British Patent 255, 831.  $P^I = P^{II} = P^{III} = P^{IV}$ .

5H by French Patent 2,315,872; 5I, 5II, 5III, 5IV

On By French Patent 2,215,973; R = R' = R'' = H, R''' = tertiary

butyl having spasmolytic properties, by British Patent 923,942;

$R^I = R^{II} = R^{III} = H$ ,  $R^{IV} = CH_3$  having spasmodic properties, by US


40 Patent 2,446,522.

**BAD ORIGINAL**

-3-


0000322

South African Patent 52/4154 discloses i.a. a compound having the formula



10 claimed to have a therapeutic utility especially as an anti-tussive.


In J. Med. Chem. 14 161 (1971) a compound of the formula




is disclosed as an antidepressant.

#### General Outline of the Invention

According to the present invention it has been found that  
25 compounds of the general formula below have advantageous, therapeutic properties:




In these compounds the dotted line represents an optional double bond. These compounds may be divided into two groups  
35 defined by the formulae




-4-

0000322



Ib

5 In the compounds of formula I Ar represents the group



10 wherein Y is bound in the 2-, 3-, or 4-position and represents a lower alkyl group, a lower alkoxy group, a halogen, a trifluoromethyl group, or an amino or a mono- or di-lower alkyl amino group, or Ar represents a pyridyl group bound in the 2-, 3- or 4-position, X represents hydrogen, a lower alkyl group,  
 15 a lower alkoxy group, a halogen, a trifluoromethyl group, an amino group or a mono- or di-lower alkyl amino group, R is a lower alkyl group and R¹ is hydrogen or a lower alkyl group. By lower alkyl and alkoxy groups are meant groups comprising up to 3 carbon atoms. Halogen may be any of the elements F,  
 20 Cl, Br or I. Therapeutically acceptable salts and bioprecursors of the compounds of the invention as well as differently hydrated or anhydrous forms of such compounds or salts are within the scope of the invention. By a bioprecursor of a therapeutically active compound is meant a compound which is  
 25 structurally different from the therapeutically active compound but which on administration to an animal or human is converted in the body to the therapeutically active compound.

The compounds of formula Ia above wherein Ar is identical to  
 30 the group




35 contain one asymmetric carbon atom. The remaining compounds of formula Ia contain two asymmetric carbon atoms and can therefore exist in two diastereomeric forms which can be separated by methods known in the art. Further the compounds of formula Ia above may be resolved into their optical

-5-

0000322

enantiomers by using optically active acids such as i.a. tartaric acid, mandelic acid, dibenzoyl tartaric acid as known in the art. The compounds of the invention may be used as mixtures of diastereomeric forms or as racemic mixtures of the pure diastereomers or as the pure enantiomers mentioned above. The therapeutic properties may reside to a greater or lesser extent in one of the enantiomers or mixtures mentioned above.

10 Due to the lack of free rotation in the double bond the compound of formula Ib in which the group Ar is not identical with the group




15

may exist in different stereoisomeric forms, that is in cis-trans isomers or, according to the IUPAC nomenclature (J. Org. Chem. 35, 2849-2867, September 1970), in an E-form and a Z-form. The compound may be used therapeutically as a mixture of geometrical isomers or in pure E or Z form. The pure geometrical isomers may be prepared from an isomer mixture, from an isomer-pure starting material or directly by a stereoselective synthesis.

25


It should be noted that in the IUPAC nomenclature compounds of formula Ib in the form of pure geometrical isomers which are similar in structure may be named the E-form for one subgroup of compounds and the Z-form for another subgroup. The two structural formulas below illustrate this fact.



35

40

E-form



Z-form

-6-

0000322

All the compounds of formula Ib further contain one asymmetric carbon atom. The compounds of formula Ib may be resolved into their optical enantiomers by using optically active acids such as i.a. tartaric acid, mandelic acid, dibenzoyl tartaric acid  
5 as known in the art. The compounds of formula Ib may be used as mixtures especially racemic mixtures, or as the pure enantiomers of the geometrical isomers mentioned above. The therapeutic properties may reside to a greater or lesser extent in one of the enantiomers or mixtures mentioned above.


10

The compounds of the invention show an activity in the central nervous system which makes them useful as neuropharmacological agents for treatment of various diseases in animals including man. The compounds are expected to be especially useful as  
15 anti-depressive, anxiolytic or tranquilizing agents in man.

#### Preferred Embodiment of the Invention

Of the compounds of the invention defined by formula I above those wherein Ar is

20



are to be specially mentioned. Of those the compounds wherein  
25 X is hydrogen are preferred and especially those wherein Y is F, Br or  $\text{CH}_3\text{O}-$ .

As preferred individual compounds should be mentioned:

30 ( $\beta$ )-3-(4-fluorophenyl)-1-methyl-3-phenylpropylamine,  
( $\beta$ )-3-(4-bromophenyl)-1-methyl-3-phenylpropylamine,  
( $\alpha$ )-3-(4-methoxyphenyl)-1-methyl-3-phenylpropylamine, and  
( $\beta$ )-3-(3-bromophenyl)-1-methyl-3-phenylpropylamine  
being non-selective inhibitors of neuronal norepinephrine and  
35 5-hydroxytryptamine uptake;

( $\alpha$ )-3-(2-bromophenyl)-1-methyl-3-phenylpropylamine,  
(E)-3-amino-1-(3-bromophenyl)-1-phenylbutene, and  
(Z)-3-amino-1-(3-bromophenyl)-1-phenylbutene



-7-

0000322

being selective inhibitors of neuronal noradrenaline uptake;  
and


- 3-amino-1,1-di-(4-methoxyphenyl)-1-butene,  
5 3,3-di-(4-fluorophenyl)-1-methylpropylamine,  
( $\beta$ )-3-(4-methoxyphenyl)-1-methyl-3-phenylpropylamine, and  
(E)-3-amino-1-(4-bromophenyl)-1-phenylbutene,  
being selective inhibitors of neuronal 5-hydroxytryptamine  
uptake; as well as salts and precursors of said compounds.  
10 The neuronal uptake mechanism are discussed further in the  
chapter "Pharmacological evaluation" below.

Generally preferred in all classes of the compounds of the  
invention are those wherein  $R^1$  represents hydrogen and R re-  
15 presents a methyl group.

#### Methods of Preparation


The compounds of the invention may be prepared by

- 20 a) reducing a compound of the formula



- wherein Ar, X and R have the meaning defined above and  $R'$  is  
25 a hydrogen atom or an alkyl, an acyl or an alkylsulfonyl  
group having 1-3 carbon atoms, to the obtention of a compound  
of formula Ia in which  $R^1$  is hydrogen, and if desired con-  
verting this primary amine to a secondary amine in a manner  
known in the art. The reduction may be carried out by known  
30 methods e.g. employing a hydride reagent such as lithium  
aluminium hydride,

- b) preparing a reactive ester of an alcohol of the formula



- wherein Ar, X and R have the meaning defined above and re-  
acting the ester obtained with an amine of the formula  $NH_2R^1$ ,  
wherein  $R^1$  has the meaning defined above. The reactive ester


-8-

0000322

may be obtained by treating the alcohol with a halogenating agent such as thionyl chloride, thionyl bromide or phosphorus tribromide, or with an arylsulphonyl halide such as p-toluene-sulphonyl chloride,

5


c) reducing a compound of the formula



Ib

10 wherein Ar, X, R and R<sup>1</sup> have the meaning defined above. The reduction may be carried out by methods known in the art e.g. by catalytical hydrogenation using catalysts such as Raney nickel, palladium on charcoal, platinum dioxide or rhodium,

15 d) reacting a ketone of the formula



IV

wherein Ar, X and R have the meaning defined above, with  
20 ammoniumformate or methylammoniumformate according to Leuckart-Wallach. The formate may be added as such, or obtained by formation in situ from formamide or methyl formamide, or from formic acid and ammonia or methylamine.


25 The intermediates of formulae II, III, and IV above are novel.

The intermediates of formula II may be prepared by reacting the ketone of the formula IV above with a hydroxylamine derivative of formula NH<sub>2</sub>OR<sup>1</sup>, wherein R<sup>1</sup> has the meaning de-  
30 fined above.

The intermediate of formula III may be prepared by hydride reduction of the compound of formula IV which, in turn, may be obtained by

35

1) reacting an alpha, beta-unsaturated ketone of the formula




V

-9-

0000322

wherein X and R have the meaning defined above, with a metal-organic reagent, such as magnesium, lithium or sodium derivative of an arylhalide of the formula Ar-Y', wherein Ar has the meaning defined above and Y' is a chlorine, bromine or iodine atom, in the presence of catalytic amounts of cuprous ions,

2) reacting a diarylcarbinol of the formula



VI


wherein Ar and X have the meaning defined above, first with thionylchloride, then with ethyl acetoacetate in the presence of suitable condensation catalyst such as sodium acetate or the like.

The compounds of formula Ia are preferably prepared by method a). The reaction according to method a) is preferably performed in diethyl ether with a slight excess of lithium aluminum hydride under inert atmosphere.

The new compounds of formula Ia may be used therapeutically as the racemic mixtures of (+)- and (-)-forms, which in the usual case are obtained at the synthesis. Isomer mixtures obtained may be resolved by methods known per se into corresponding optically active modifications. If desired, the optically active modification may be prepared by way of direct synthesis, e.g. via an optically active compound as described above.

The compounds of the formula I in the invention may be prepared by:

e) Dehydration of a carbinol of the formula



VII

to a compound of formula Ib.

BAD ORIGINAL


-10-

0000322

The dehydration of the starting material may be done by means of treatment with hydrochloric acid HCl and heating of the reacting mixture. The dehydration of the starting material may also be done by means of other types of acid-catalysis, such as by means of sulfuric acid H<sub>2</sub>SO<sub>4</sub>, phosphoric acid H<sub>3</sub>PO<sub>4</sub>, potassium hydrogen sulphate KHSO<sub>4</sub>, or oxalic acid (COOH)<sub>2</sub>. Other methods for the dehydration of the starting material to the formation of a compound of the formula I are dehydration using phosphoroxichloride POCl<sub>3</sub> in pyridine, and dehydration with thionylchloride, SOCl<sub>2</sub>, in pyridine. Also a catalytic dehydration of the starting material may be used. The dehydration is in this case carried out at a temperature of about 300 to 500°C using a catalyst such as kaolin, alumina or aluminium oxide.

15

f) Reaction of a compound of the formula




VIII

wherein Z is a leaving group such as F, Cl, Br, I, OSO<sub>2</sub>R<sup>1</sup>, wherein R<sup>1</sup> is alkyl, aralkyl or aryl, with an amine of the formula NH<sub>2</sub>R<sup>1</sup> or with a derivative thereof such as hexamethylenetetraamine, alkaliphtalimide, lithium bisbenzenesulfeneamide, guanidine, sodium cyanate, sodium azide, a carboxamide or a sulfonamide. When an amine derivative is alkylated the product obtained is subsequently hydrolyzed or in some other way converted into a primary or secondary amine of formula Ib. A preferred amine derivative is potassium phtalimide.

30

This reaction is also useful for preparation of the compounds of formula Ia by employing as a starting material a saturated compound corresponding to the compound of formula VIII.

g) Oxidation (halogenation) of the benzylic carbon atom of a compound of formula IX




IX

-11-


0000322

e.g. with N-bromosuccinimide, R'', R''' being protective groups for the amino function preferably joined as the group



5

or alternatively corresponding to one of the amine derivatives specified under f) above, followed by elimination of the group formed by oxidation at the benzyl carbon atom, giving a compound of the formula




X

which is transformed into the compound of formula Ib by splitting off the groups R'' and R''' , e.g. by hydrolysis or hydrazinolysis. In this manner it is possible to synthesize the compounds of formula Ib from the corresponding saturated compounds of formula Ia, by introducing in those compounds protective groups R'' and R''' in a known manner to the content of a compound of formula IX.

The intermediates of formulae VII, VIII, IX and X above are novel.


The intermediate of formula VII may be obtained by preparing a Grignard or lithium compound from a halobenzene of the formula



Hal = Br, Cl

30

and reacting it with an ester of the formula



wherein R'' is an alkyl, aralkyl or aryl group.


35

The intermediate of formula VIII may be isolated (cf. Example 15) and subsequently dehydrated or alternatively the crude product in the preparation of formula VII may be dehydrated directly (cf. Example 17).

-12-

0000322

The intermediate of formula VIII may be obtained from a compound of formula



wherein X, Ar and R have the meaning defined above, by oxidation of the allylic carbon to the formation of a compound of the formula

10



wherein  $Z^1$  represents Z or OH, e.g. by reaction with N-bromo-  
15 succinimide ( $Z^1 = \text{Br}$ ) or selenium dioxide ( $Z^1 = \text{OH}$ ). The latter compound may be transformed to a reactive derivative III by treatment with an agent such as  $\text{SOCl}_2$ ,  $\text{SOBr}_2$  or  $\text{PBr}_3$  or with  $\text{ClSO}_2\text{R}^1$ .

## 20 Pharmaceutical Preparations

In clinical practice the compounds of the present invention will normally be administered orally, rectally or by injection, in the form of pharmaceutical preparations comprising the active ingredient either as a free base or as a pharmaceutically  
25 acceptable non-toxic, acid addition salt, e.g. the hydrochloride; hydrobromide, lactate, acetate, phosphate, sulphate, sulphamate, citrate, tartrate, oxalate and the like in association with a pharmaceutically acceptable carrier. Accordingly, terms relating to the novel compounds of this invention whether  
30 generically or specifically are intended to include both the free amine base and the acid addition salts of the free base, unless the context in which such terms are used, e.g. in the specific examples would be inconsistent with the broad concept. The carrier may be a solid, semisolid or liquid diluent or  
35 capsule. These pharmaceutical preparations constitute a further aspect of this invention. Usually the active substance will constitute from 0.1 to 99 % by weight of the preparation, more specifically between 0.5 and 20 % by weight for preparations intended for injection and between 2 and 50 % by

-13-

0000322

weight for preparations suitable for oral administration.

To produce pharmaceutical preparations containing a compound of the invention in the form of dosage units for oral application the selected compound may be mixed with a solid pulverulent carrier, e.g. lactose, saccharose, sorbitol, mannitol, starches such as potato starch, corn starch or amylopectin, cellulose derivatives, a binder such as gelatine or polyvinylpyrrolidone, and a lubricant such as magnesium stearate, calcium stearate, polyethylene glycol waxes, and the like, and then compressed to form tablets. If coated tablets are required, the cores, prepared as described above, may be coated with a concentrated sugar solution which may contain, e.g. gum arabic, gelatine, talcum, titanium dioxide, and the like. Alternatively, the tablet can be coated with a lacquer in a readily volatile organic solvent or mixture of organic solvents. Dyestuff may be added to these coatings in order to readily distinguish between tablets containing different active substances or different amounts of the active compounds.

For the preparation of soft gelatine capsules (pearlshaped closed capsules) consisting of gelatine and for example, glycerol or similar closed capsules, the active substance may be admixed with a vegetable oil. Hard gelatine capsules may contain granulates of the active substance in combination with solid, pulverulent carriers such as lactose, saccharose, sorbitol, mannitol, starches (e.g. potato starch, corn starch or amylopectin), cellulose derivatives or gelatine.

Dosage units for rectal application can be prepared in the form of suppositories comprising the active substance in admixture with a neutral fatty base, or gelatine rectal capsules comprising the active substance in admixture with vegetable oil or paraffin oil.

Liquid preparations for oral application may be in the form of syrups or suspensions, for example, solutions containing from about 0.2 % to about 20 % by weight of the active substance herein described the balance being sugar and a mixture

-14-

0000322

of ethanol, water, glycerol, and propyleneglycol. Optionally such liquid preparations may contain colouring agents, flavouring agents, saccharine and carboxymethylcellulose as a thickening agent.

5

Solutions for parenteral applications by injection can be prepared in an aqueous solution of a water-soluble pharmaceutically acceptable salt of the active substance preferably in a concentration of from about 0.5 % to about 10 % by weight. These  
10 solutions may also contain stabilizing agents and/or buffering agents and may conveniently be provided in various dosage unit ampoules.

Suitable daily doses of the compounds of the invention in  
15 therapeutic treatment is 25 to 250 mg for peroral administration, preferably 50 to 150 mg, and 5 to 50 mg for parenteral administration preferably 10 to 30 mg. A preparation in dosage unit form for oral administration may contain 10 to 50 mg, preferably 10 to 25 mg of active substance per dosage unit.

20

#### Working Examples

##### Example 1. Preparation of 4-(3-bromophenyl)-4-phenylbutan-2-one

To 2.4 g (0.10 mol) of magnesium turnings, covered with 20 ml  
25 of anhydrous diethyl ether and treated with some crystals of iodine under nitrogen atmosphere, 23.5 g (0.10 mol) of 1,3-dibromobenzene in 30 ml of ether was added. The rate of addition was adjusted, to maintain a gentle reflux of the solvent. When the Mg turnings had disappeared (30 min) 0.75 g  
30 (0.005 mol) of cuprous bromide was added and the mixture was stirred for 10 min at room temperature. A solution of 13.6 g (0.09 mol) of benzalacetone in 100 ml of ether was added dropwise at +10°C. Then the reaction mixture was allowed to reach room temperature for 2 hours. The mixture was poured into  
35 400 ml of 10 % aqueous solution of ammonium chloride, the aqueous layer was separated and the product was extracted with 2 x 200 ml of ether. The ethereal layer was washed with water, dried with Na<sub>2</sub>SO<sub>4</sub> and the solvent was evaporated. The crude 3-(3-bromophenyl)-3-phenylbutan-2-one obtained (20.5 g,



-15-

0000322

0.086 mol) was added to a solution of 20 g (0.28 mol) of hydroxylamine hydrochloride in 300 ml of ethanol and 100 ml of anhydrous pyridine. The mixture was heated under reflux for 4 hours. After cooling the solvent was evaporated in vacuo and the ketoxime was extracted with ether from an aqueous solution. Washing of the extract with water and drying followed by evaporation of the solvent gave 27.5 g of 3-(3-bromophenyl)butan-2-one oxime.

10 Example 2. Preparation of 3-(3-bromophenyl)-1-methyl-3-phenyl-propylamine oxalate. (Method a)

To the oxime obtained according to Example 1 (27.5 g) 100 ml of carbon tetrachloride was added and evaporated twice in order to remove traces of water. The residue was dissolved in a mixture of 300 ml of anhydrous ether and 150 ml of anhydrous tetrahydrofuran. To the stirred mixture 3.5 g (0.0086 mol) of lithium aluminium hydride was added in portions under nitrogen atmosphere at room temperature. The reaction mixture was stirred for 8 hours. Then 25 ml of 2 M NaOH was dropwise added and the precipitated inorganic salts were removed by filtration. The filtrate was shaken with 3 x 300 ml of 1 M HCl and the combined aqueous layers were made alkaline by addition of 35 ml of 30 % NaOH. Extraction with 3 x 200 ml of methylene chloride, washing, drying and evaporation of the solvent gave 8.9 g of the primary amine as an oil. To a hot solution of 7.9 g (0.026 mol) of the amine in 100 ml of isopropylalcohol 1.1 g (0.014 mol) of oxalic acid in 10 ml of ethanol was added. 6.2 g of the diamine oxalate was collected. Recrystallization from 160 ml of a mixture of ethanol and isopropylalcohol (1:1) yielded 4.43 g, mp 134-138°C.

Analysis: C calcd 58.5 %, found 58.7 %; H calcd 5.48 %, found 5.80 %; N calcd 4.01 %, found 4.07 %.

35 Example 3. Separation of 3-(3-bromophenyl)-1-methyl-3-phenyl-propylamine into its diastereomers

The free amine (2.9 g, 0.009 mol) obtained from 3.3 g of the oxalate prepared according to Example 2, was dissolved in 40 ml of ethyl acetate. A hot solution of 1.1 g (0.009 mol) of

-16-

0000322

maleic acid in 20 ml of ethanol was added. There was obtained 1.3 g of the maleate. Recrystallization from 18 ml of isopropylalcohol gave 0.65 g of the pure alpha isomer, mp 163-165°C. The high field part of the NMR spectrum (CCl<sub>3</sub>) displayed a triplet at 4.1 ppm (J=7.8 Hz), a quartet at 2.8 ppm (J=6.2 Hz), a doublet at 2.0 ppm (two protons) and a doublet at 1.5 ppm (J=6.1 Hz) (three protons).

Analysis: C calcd 57.15 %, found 57.45 %; H calcd 5.28 %, found 5.35 %; Br calcd 19.01 %, found 19.05 %; N calcd 3.33 %, found 3.20 %; O calcd 15.23 %, found 15.00 %.

The solvents of the first mother liquors of the diamine oxalate prepared according to example 2 were evaporated and the residue was extracted with ether from an alkaline solution. There was obtained 1.2 g of free amine. The fumarate was prepared in ethyl acetate from half an equivalent of fumaric acid and recrystallized twice from acetonitrile-isopropylalcohol affording 0.25 g of the pure beta isomer as the diamine fumarate, mp 184-186°C.

Analysis: C calcd 59.7 %, found 60.3 %; H calcd 5.6 %, found 5.7 %; N calcd 3.9 %, found 3.7 %.

#### 25 Example 4

(α)-3-(4-fluorophenyl)-1-methyl-3-phenylpropylamine oxalate, mp 186-189°C (EtOH-EtOAc, 1:1) and (β)-3-(4-fluorophenyl)-1-methyl-3-phenylpropylamine hydrochloride, mp 171-172°C (EtOAc) were prepared from 3-(3-fluorophenyl)-3-phenylbutan-2-one oxime in accordance with Examples 2 and 3.

#### Example 5

(α)-3-(4-bromophenyl)-1-methyl-3-phenylpropylamine maleate, mp 168-170°C (EtOH-EtOAc, 1:1) and (β)-3-(4-bromophenyl)-1-methyl-3-phenylpropylamine maleate, mp 161-162°C (i-PrOH-EtOAc, 3:1) were prepared from 3-(4-bromophenyl)-3-phenylbutan-2-one oxime in accordance with Examples 2 and 3.

-17-

0000322

Example 6

( $\alpha$ )-3-(4-methoxyphenyl)-1-methyl-3-phenylpropylamine maleate, mp 146-148°C (i-PrOH) and ( $\beta$ )-3-(4-methoxyphenyl)-1-methyl-3-phenylpropylamine maleate, mp 124-133°C (EtOAc) were prepared from 3-(4-methoxyphenyl)-3-phenylbutan-2-one oxime in accordance with Examples 2 and 3.

Example 7

( $\alpha$ )-1-methyl-3-(4-trifluoromethylphenyl)-3-phenylpropylamine maleate, mp 165-166°C and ( $\beta$ )-1-methyl-3-(4-trifluoromethylphenyl)-3-phenylpropylamine maleate, mp 165-167°C were prepared from 3-(4-trifluoromethylphenyl)-3-phenylbutan-2-one oxime in accordance with Examples 2 and 3.

15 Example 8. Preparation of 4-(2-bromophenyl)-4-phenylbutan-2-one

To a solution of 24.3 g (0.13 mol) of 2-bromobenzaldehyde in 60 ml of acetone, 1.0 ml of 10 M NaOH was slowly added at 0°C. The reaction mixture was allowed to reach room temperature and stirred for another 2 hours. Then it was poured into 400 ml of water, to which 10 ml of 2 M HCl had been added. Extraction with ether, drying and evaporation of the solvent gave 13.9 g of 2-bromobenzalacetone as an oil. This was dissolved in 150 ml of ether and added to a Grignard reagent prepared from 1.1 g (0.045 mol) of magnesium turnings 6.6 g (0.042 mol) of bromobenzene and 0.2 g of CuBr in 150 ml of ether. The mixture was stirred under nitrogen atmosphere for 2 hours. It was poured into 450 ml of ice-water to which 18 g of ammonium chloride had been added. Extraction with ether gave 11.0 g of the desired ketone as an oil.

30

Example 9

( $\alpha$ )-3-(2-bromophenyl)-1-methyl-3-phenylpropylamine maleate, mp 145-146°C (i-PrOH) and ( $\beta$ )-3-(2-bromophenyl)-1-methyl-3-phenylpropylamine maleate, mp 135-137°C (EtOH-EtOAc, 1:4) were prepared from 3-(2-bromophenyl)-3-phenylbutan-2-one oxime in accordance with Examples 2 and 3.

-18-

0000322

Example 10. Preparation of N,1-dimethyl-3-(4-bromophenyl)-3-phenylpropylamine

The free base of 3-(4-bromophenyl)-1-methyl-3-phenylpropylamine (0.07 g, 0.0023 mol) was dissolved in 50 ml of chloroform. 1.2 ml (0.0024 mol) of 2 M NaOH and 0.25 g (0.0024 mol) of ethyl chloroformate were added separately and dropwise with vigorous stirring at 15°C. Stirring was continued for 2 hours at room temperature.

- 10 Then 25 ml of water was added and the organic layer was separated, dried, and the solvent was evaporated to give 1.0 g of N-ethoxycarbonyl-3-(4-bromophenyl)-1-methyl-3-phenylpropylamine as a colourless oil.
- 15 Treatment of the carbamate with 0.25 g (0.006 mol) of  $\text{LiAlH}_4$  in 60 ml of ether for 14 hours gave 0.20 g of the secondary amine after extraction with ether/hydrochloride and  $\text{NaOH}$ /ether. The hydrochloride was prepared and recrystallized from 18 ml of acetone to give 0.12 g, mp 146-148°C.

20

The oxalate had mp 106-111°C from acetone.

Example 11. Preparation of 3-(3-bromophenyl)-1-methyl-3-phenylpropylamine. (Method d)

- 25 A mixture of 22.1 g (0.073 mol) of 4-(3-bromophenyl)-4-phenylbutan-2-one and 230 ml of formamide was heated for 8 hours at 180°C. After cooling water was added and the product was taken up in ether. Drying and evaporation of the solvent gave 30.5 g of a residue.

30

- To this residue 35 ml of conc. hydrochloric acid was added and the mixture was heated under refluxing conditions for 3 hours. Water was added and the non basic materials were removed by shaking the reaction mixture with 120 ml of ether.
- 35 The aqueous layer was separated and made alkaline by addition of 120 ml of 10 N NaOH. Extraction with 3 x 200 ml of ether, drying ( $\text{Na}_2\text{SO}_4$ ) and evaporation of the solvent gave 12.9 g of the desired amine.

-19-

0000322

The maleate was prepared by addition of a hot ethanolic solution of 4.9 g of maleic acid into a warm solution of the amine in 100 ml of ethyl acetate. Recrystallization from EtOH-EtOAc gave 8.4 g of the maleate, mp 153-161°C.

5

Analysis: C calcd 57.2 %, found 57.5 %; H calcd 5.28 %, found 5.35 %; Br calcd 19.0 %, found 19.1 %; N calcd 3.33 %, found 3.20 %; O calcd 15.2 %, found 15.0 %.

10 Example 12. Preparation of 3,3-di-(4-fluorophenyl)-1-methyl-propylamine hydrochloride (Method c)

3,3-di-(4-fluorophenyl)-1-methylallylamine (0.9 g, 0.003 mol) was dissolved in 100 ml of ethanol and transferred to a Parr hydrogenation flask. 1.0 ml of concentrated hydrochloric acid was added followed by 0.2 g of 5 % palladium on charcoal. The hydrogenation was effectuated at a pressure of 3.5 atm for 5.5 hours. The reaction mixture was filtered to remove the catalyst and the solvent of the filtrate was evaporated. Crystallization from EtOAc-i-Pr<sub>2</sub>O gave 0.75 g of the desired product, mp 225-229°C.

25

Analysis: C calcd 64.5 %, found 64.5 %; H calcd 5.09 %, found 5.11 %; Cl calcd 11.9 %, found 11.8 %; F calcd 12.8 %, found 12.7 %; N calcd 4.70 %, found 4.55 %.

By the same method there were prepared from the appropriate allylamines:

Example 13

30 By the method of Example 12 3,3-di-(4-methoxyphenyl)-1-methyl-propylamine fumarate, mp 153-157°C, (EtOH-i-Pr<sub>2</sub>O) was prepared from the corresponding allylamine.

Example 14

35 By the method of Example 12 1,3', 3"-trimethyl-3,3-diphenyl-propylamine oxalate, mp 214-215°C, (EtOH-EtOAc) was prepared from the corresponding allylamine.

-20-

0000322

Example 15. Preparation of 4,4-di-(4-bromophenyl)-4-hydroxy-2-butylamine oxalate

A solution of 81.5 g (0.35 mol) 1,4-dibromobenzene in 500 ml of diethyl ether was added to a stirred mixture of 8.3 g (0.35 mol) magnesium turnings in 25 ml diethyl ether at such a rate that reflux was maintained. After an additional stirring for 1.5 hours at room temperature the mixture was cooled in an ice-bath and a solution of 11.3 g (0.11 mol) of ethyl 3-aminobutyrate in 25 ml of diethyl ether was added during 15 min.

10

The mixture was stirred for 1.25 hours at ice-cooling and then for 2.5 hours at reflux. An aqueous cold solution of 25 g ammonium chloride was slowly added, and after stirring the mixture was extracted twice with ether. The ethereal layer was dried over sodium sulphate and the amine was precipitated (10.9 g, 20 % yield) with oxalic acid dissolved in ether. M.p. 191-194°C.

Elemental analysis:  $C_{18}H_{19}Br_2NO_5$ : Found: C 44.5, H 4.0, N 2.7, 20 and O 16.6 %. Calculated: C 44.20, H 3.91, N 2.86 and O 16.35 %.

Example 16. Preparation of 3-amino-1,1-di(4-bromophenyl)-1-butene hydrochloride (Method a)

A solution of 3.8 g 4,4-di(4-bromophenyl)-4-hydroxy-2-butyl-25 amine oxalate, 25 ml acetic acid and 5 ml conc. aqueous hydrogen chloride was heated under reflux for 30 min. The solvent was evaporated and the residue was made alkaline with sodium hydroxide and extracted twice with ether. The ethereal layer was dried over sodium sulphate and the solvent was evaporated. 30 Acetonitrile and hydrogen chloride in ether were added and the hydrochloride of the title compound (1.7 g) was obtained after recrystallization from acetonitrile/ether. M.p. 216-220°C.

35 Elemental analysis:  $C_{16}H_{15}Br_2ClN$ ; Found: C 46.7, H 3.9, Cl 8.8 and N 3.1 %. Calculated: C 46.02, H 3.36, Cl 8.49 and N 3.35 %.

-21-

0000322

Example 17. Preparation of 3-amino-1,1-di(4-methoxyphenyl)-  
-1-butene fumarate (Method e)

A solution of 58.0 g (0.31 mol) of 4-bromoanisole in 300 ml diethyl ether was added dropwise to a stirred mixture of 7.78 g (0.32 mol) magnesium turnings in 250 ml of diethyl ether during 2 hours. The mixture was stirred at room temperature for another 1.5 hours and then heated under reflux for 1.5 hours. The mixture was cooled in an ice-bath and a solution of 10.3 g (0.10 mol) ethyl 3-aminobutyrate in 25 ml diethyl ether was added during 20 min. After stirring over night at room temperature an aqueous solution of 16.6 g (0.31 mol) ammonium chloride was slowly added. The mixture was stirred and then made alkaline and filtered. After separation of the ether phase the aqueous phase was extracted with ether. The combined ethereal layers were extracted twice with 2 M hydrogen chloride. The aqueous phase was made alkaline and extracted with ether and dried over sodium sulphate. After evaporation of the solvent 10.1 g of yellow oil was obtained, which was dissolved in 75 ml of acetic acid and 15 ml of conc. aqueous hydrogen chloride.

The solution was heated under reflux for 30 min and then the solvent was evaporated. The residue was made alkaline and extracted with ether. The ethereal layer was washed with water and extracted with 0.5 M hydrogen chloride. The aqueous phase was washed with ether, made alkaline and extracted with ether. After drying over sodium sulphate the ether was evaporated to give 5.9 g of the title compound as an oil in 21 % yield.

The amine was converted to the fumarate, which was recrystallized twice from ethanol/ethyl acetate/hexane to give 6.2 g (17 % yield) of the fumaric acid salt in the form  $C_{18}H_{21}NO_2 \cdot 3/4C_4H_4O_4$ . M.p. 167-167.5°C.

Elemental analysis:  $C_{21}H_{24}NO_5$ : Found: C 57.8, H 6.52, N 3.89 and O 21.5 %. Calculated: C 58.09, H 6.53, N 3.78 and O 21.60 %.

-22-

0000322

Example 18

3-amino-1,1-di-(4-fluorophenyl)-1-butene fumarate was prepared according to Example 17. M.p. 225-231°C.

5 Example 19

3-amino-1,1-di-(3-methylphenyl)-1-butene hydrochloride was prepared according to Example 17. M.p. 216-217.5°C.

Example 20. Preparation of 1-(4-bromophenyl)-1-phenyl-butene

10 Sodium dimethylsulfoxide in DMSO, prepared by heating 3.8 g (0.08 mol) sodium hydride (50 % in oil) in 100 ml dimethylsulfoxide at 80°C for 40 min, was mixed with 27.0 g (0.07 mol) propyltriphenylphosphonium bromide, prepared by heating propylbromide and triphenylphosphine in toluene at reflux temperature  
15 for 14 hours. The mixture was stirred under nitrogen atmosphere at room temperature for 1.5 hours. Then a solution of 13.1 g (0.05 mol) of 4-bromobenzophenone, in a mixture of 100 ml dimethylsulfoxide and 100 ml of anhydrous tetrahydrofuran was added at room temperature. The reaction mixture was  
20 stirred for 2 hours, then it was poured into 1.000 ml of ice-water. The product was extracted with 3 x 300 ml of ether, the combined ethereal layer was washed with 100 ml of water. Drying (Na<sub>2</sub>SO<sub>4</sub>) and evaporation of the solvent gave 18 g of oily residue. After trituration with 100 ml of diisopropylether crystals  
25 of triphenylphosphine oxide was separated by filtration. Distillation of the filtrate at 4 Pa gave 12.2 g of 1-(4-bromophenyl)-1-phenyl-butene, b.p. 120-130°C. Yield 85 %.

Example 21. Preparation of 3-(4-bromophenyl)-3-phenyl-1-30 -methylallylbromide

To a solution of 22.6 g (0.0787 mol) of 1-(4-bromophenyl)-1-phenylbutene in 800 ml of carbon tetrachloride 14.0 g (0.0787 mol) of N-bromosuccinimide was added. The mixture was heated to reflux temperature after the addition of 0.7 g  
35 of alpha, alpha'-azobiscytronitrile. After 3.5 hours all NBS had been consumed and the reaction mixture was cooled and filtered. 7.8 g of succinimide was separated and the volume of the filtrate was reduced to 50 ml by evaporation at 35°C. TLC of a sample on silica in diisopropylether-

BAD ORIGINAL



-23-

0000322

hexane (1:1) showed a new spot at  $R_f$  0.62 (the starting olefin had  $R_f$  0.55). The material was being used without further isolation or purification due to its high reactivity with nucleophilic agents.

5

Example 22. Preparation of 3-(N-phthalimido)-1-(4-bromophenyl)-1-phenylbutene (Method f)

A solution of crude 3-(4-bromophenyl)-3-phenyl-1-methylallyl-bromide (29 g, 0.08 mol) in 50 ml carbon tetrachloride was  
10 mixed with 15.0 g (0.08 mol) of N-potassiumphthalimide and 120 ml of anhydrous dimethylformamide. The mixture was stirred at 50°C for 14 hours.

Dilution with water (excess) and extraction of the product with  
15 diethylether gave 19 g of an oil after drying and evaporation of the solvent. Column chromatography on silica with diisopropylether as the eluent afforded the geometrical isomers: 2.6 g of the (Z)-form,  $R_f$  = 0.30, and 3.7 g of the (E)-form,  $R_f$  = 0.26 in diisopropylether-hexane (1:1).

20

Example 23. Preparation of (Z)-3-amino-1-(4-bromophenyl)-1-phenylbutene maleate

To a stirred solution of 0.43 g (0.001 mol) of (Z)-3-phthalimido-1-(4-bromophenyl)-1-phenylbutene in 30 ml of methanol  
25 0.25 g (0.005 mol) of hydrazine hydrate was added at room temperature. In order to dissolve all the phthalimide 10 ml of carbon tetrachloride was added. The mixture was stirred and heated at 60°C for 2 hours. After cooling the solvent was removed in vacuo and the residue was taken up in ether.  
30 The product was extracted with 3 x 50 ml of 0.5 M HCl, the combined aqueous layer was made alkaline with 10 M NaOH and extracted with 2 x 50 ml of ether. Drying and evaporation of the solvent gave 0.23 g of the title compound as an oil. The maleate had m.p. 174-176°C from ethanol. The UV spectrum  
35 in ethanol had  $\lambda_{max}$  237 nm.

Elemental analysis:  $C_{20}H_{20}BrNO_4$ ; Found: C 57.1, H 4.80, Br 20.3, N 3.10 and O 15.2 %. Calculated: C 57.43, H 4.82, Br 19.10, N 3.35 and O 15.30 %.

-24-

0000322

Example 24

(E)-3-amino-1-(4-bromophenyl)-1-phenylbutene oxalate was prepared from the corresponding phthalimide according to Example 23. M.p. 145-148°C.

5

Example 25. Preparation of (Z)-3-amino-1-(3-bromophenyl)-1-phenylbut-1-ene maleate (Method f)

To a stirred solution of 0.43 g (0.001 mol) of (Z)-3-phthalimido-1-(3-bromophenyl)-1-phenylbutene in 40 ml of methanol, 10 0.35 g (0.007 mol) of hydrazine hydrate was added at room temperature. The mixture was heated under reflux for 2.5 hours. The solvent was evaporated and the residue was taken up in ether. Extraction with 3 x 25 ml of 0.5 M HCl followed by alkalization of the combined aqueous layer with 10 M NaOH 15 and extraction with 2 x 50 ml of ether gave 0.19 g of residue after drying and evaporation of the solvent. The maleate was prepared from 15 ml ethylacetate - ethanol (2:1) to give 0.12 g (28 %). M.p. 198-200°C.

20 Elemental analysis:  $C_{20}H_{20}BrNO_4$ ; Found: C 56.3, H 4.7, and N 3.2 %. Calculated: C 57.43, H 4.92 and N 3.35 %.

Example 26

(E)-3-amino-1-(3-bromophenyl)-1-phenylbutene hydrochloride was 25 prepared from the corresponding phthalimide according to Example 25. M.p. 118-123°C.

Example 27

(Z)-3-amino-1-(4-bromophenyl)-1-(3-pyridyl)-butene oxalate was 30 prepared from the corresponding phthalimide according to Example 25. M.p.

Example 28. Preparation of 3-(N-Phthalimido)-1-(4-bromophenyl)-1-phenylbutane

35 3-(4-bromophenyl)-1-methyl-3-phenylpropylamine as the free base (41.2 g, 0.136 mol) was dissolved in 350 ml of acetic acid. Phthalic anhydride (20.0 g, 0.135 mol) was added and the mixture was heated with stirring under reflux (bath temperature 120°C) for 2 hours. After cooling the solvent was evaporated in vacuo.

-25-

0000322

The residue was shaken with a mixture of 300 ml of ether and 500 ml of 2 M NaOH. The ethereal layer was separated and washed with 100 ml 1 M hydrochloric acid. Drying and evaporation gave 43.5 g of a tan oil. Thin layer chromatography on silica in diisopropylether showed one spot of Rf 0.42. NMR showed a four proton multiplet at 7.7 ppm from TMS, characteristic of phthalimides. The material was used without further purification. Yield: 63 %.

10 Example 29. Preparation of 3-(N-phthalimido)-1-(4-bromophenyl)-1-phenylbut-1-ene (Method g)

To a stirred solution of 24.6 g (0.057 mol) 3-(N-phthalimido)-1-(4-bromophenyl)-1-phenylbutane in 400 ml of carbon tetrachloride 10.3 g (0.057 mol) of N-bromosuccinimide was added. 15 The mixture was stirred and 0.5 g of alpha, alpha-azaisobutyronitrile was added as radical initiator. Stirring was continued under reflux temperature for 2.5 hours. The reaction mixture was cooled and filtered. Upon evaporation of the solvent 32.4 g of a residue was obtained. NMR of a 20 sample in CCl<sub>4</sub> showed a double quartet at 5.0 ppm from TMS and a doublet at 6.6 ppm. TLC on silica in diisopropylether showed a spot with Rf 0.25, identical with the Rf-value of the material prepared according to Example 22.

25 The following examples illustrate how the compound of the present invention may be included in pharmaceutical preparations.

Example 30. Preparation of soft gelatin capsules

30 500 g of active substance were mixed with 500 g of corn oil, whereupon the mixture was filled in soft gelatin capsules, each capsule containing 100 mg of the mixture (i.e. 50 mg of active substance).

35 Example 31. Preparation of soft gelatin capsules

500 g of active substance were mixed with 750 g of pea nut oil, whereupon the mixture was filled in soft gelatin capsules, each capsule containing 125 mg of the mixture (i.e. 50 mg of active substance).

-25-

0000322

Example 32. Preparation of tablets

50 kg of active substance were mixed with 20 kg of silicic acid of the trade mark Aerosil. 45 kg of potato starch and 50 kg of lactose were mixed therewith and the mixture was moistened with a starch paste prepared from 5 kg of potato starch and distilled water, whereupon the mixture was granulated through a sieve. The granulate was dried and sieved, whereupon 2 kg of magnesium stearate was mixed into it. Finally the mixture was pressed into tablets each weighing 172 mg.

10

Example 33. Preparation of an emulsion

100 g of active substance were dissolved in 2500 g of pea nut oil. From the solution thus obtained, 90 g of gum arabic, aroma and colouring agents (q.s.) and 2500 g of water an emulsion was prepared.

15

Example 34. Preparation of a syrup

100 g of active substance were dissolved in 300 g of 95 % ethanol, whereupon 300 g of glycerol, aroma and colouring agents (q.s.) and 1000 ml of water were mixed therein. A syrup was obtained.

20

Example 35. Preparation of a solution

100 g of active substance were dissolved in 2000 g of polyoxyethylene sorbitane monooleate, whereupon flavouring agents and colouring agents (q.s.) and water to 5000 ml was mixed therein. A drop solution was obtained.

25

Example 36. Preparation of effervescent tablets

100 g of active substance, 140 g of finely divided citric acid, 100 g of finely divided sodium hydrogen carbonate, 3.5 g of magnesium stearate and flavouring agents (q.s.) were mixed and the mixture was pressed into tablets each containing 100 mg of active substance.

30

Example 37. Preparation of a drop solution

100 g of active substance were mixed with 300 g of ethanol, whereupon 300 g of glycerol, water to 1000 ml, aroma and flavouring agents (q.s.) and 0.1 N sodium hydroxide solution

35

-27-

0000322

(to pH 4.5 to 5.5) was added while stirring. A drop solution was obtained.

Example 38. Preparation of a sustained release tablet

5 200 g of active substance were melted together with 50 g of stearic acid and 50 g of carnauba wax. The mixture thus obtained was cooled and ground to a particle size of at most 1 mm in diameter. The mixture thus obtained was mixed with 5 g of magnesium stearate and pressed into tablets each weighing 305 mg.  
10 Each tablet thus contains 200 mg of active substance.

Pharmacological Evaluation

Depressions are considered to be connected with changes in the biochemical processes of the brain which processes control the  
15 mood. The nature of these biochemical processes are largely unknown but in depressive states there is evidence for a decreased activity of monoaminergic brain neurons. The monoamines, noreadrenaline (NA), dopamine (DA) and 5-hydroxytryptamine (5-HT), are of great interest in this respect.

20

It has been demonstrated that NA, DA and 5-HT is localised in three different types on neurons and may function as transmitters in the central nervous system. The monoamines are stored in special structures, granules, situated in enlargements of the nerve endings, varicosities. The varicosity is  
25 separated from the effector neuron by a space, the synaptic cleft or spatium. As a result of a nerve stimulation the transmitter is released from the granule into the synaptic cleft and reaches the receptor of the effector neuron and  
30 generates a nerve impulse. After impulse generation the amines are inactivated by mainly two mechanisms: a re-uptake mechanism at the cell membrane and enzymatic conversion by catechol-O-methyltransferase to form methylated metabolites. There is also an inactivating enzyme within the varicosities,  
35 monoamine oxidase (MAO), that is stored in the mitochondria and inactivates the amines intracellularly.

When MAO-inhibitors are administered, an increased amount of transmitter substance becomes available for release at the  
40 nerve ending.

-28-

0000322

Another way of increasing the amine levels at the receptor is exerted by the tricyclic antidepressants. It has been shown that this type of compounds inhibits the re-uptake mechanism of NA and 5-HT, and the antidepressive action is assumed to be related to the uptake inhibition of NA and 5-HT.

It has been proposed, that some depressions are caused by deficiency in either one of the neurotransmitters and some of deficiency in both.

10

An antidepressant effect should thus be obtained with compounds which are able to inhibit the re-uptake of one or both NA and 5-HT.

#### 15 Pharmacological Methods

The test method described in Europ. J. Pharmacol. 17, 107, 1972. This method involves the measurement of the decrease in the uptake of  $^{14}\text{C}$ -5-hydroxytryptamine ( $^{14}\text{C}$ -5-HT) and  $^3\text{H}$ -noradrenaline ( $^3\text{H}$ -NA) in brain slices from mice after in vivo and in vitro administration of the test substance.

#### Inhibition of the uptake of $^{14}\text{C}$ -5-HT and $^3\text{H}$ -NA in vitro and in vivo

The test substances were administered intraperitoneally half an hour before the animals were killed. The midbrain was taken out, sliced and incubated in a mixture consisting of 0.2 nmole of  $^{14}\text{C}$ -5-HT, 0.2 nmole of  $^3\text{H}$ -NA and 11  $\mu\text{mole}$  of glucose in 2 ml of Krebs Henseleit-buffer, pH 7.4 per 100 mg of brain slices. The incubation time was 5 minutes with 5 minutes of preincubation before the labelled amines were added. The slices were dissolved in Soluene<sup>®</sup> and the amounts of radioactive amines taken up were determined by liquid scintillation. The doses producing 50 per cent decrease of the active uptake ( $\text{ED}_{50}$ ) of  $^{14}\text{C}$ -5-HT and  $^3\text{H}$ -NA were determined graphically from dose response curves. Active uptake is defined as that part of the radioactive uptake which is inhibited by a high concentration of cocaine.

-29-

0000322

In the vitro administration method slices of mouse midbrain were preincubated for 5 minutes with solution of the compound to be tested and then incubated as described above. The concentration producing 50 per cent inhibition of the active uptake  $IC_{50}$  of  $^{14}C$ -5-HT and 3-H-NA was determined graphically from dose response curves.

The test results are given in Table I.

TABLE I  
Inhibition of neuronal uptake of 5-hydroxytryptamine and noradrenaline in slices from mouse brain



Compound	Inhibition of the uptake in vitro			Inhibition of the uptake in vivo i.p.		
	IC <sub>50</sub> (μM)	NA	5-HT	EO <sub>50</sub> (μmole/kg)	NA	5-HT
Compounds of the invention						
	$\text{Ar} \cdot$	Y	X	R	R <sup>1</sup> isomer salt <sup>1</sup>	
CPK 163		4-F	H	CH <sub>3</sub>	H	α Ox 0.54 0.28 32
CPK 170	"	4-F	H	CH <sub>3</sub>	H	β HCl 0.25 0.18 37
CPK 185	"	2-Br	H	CH <sub>3</sub>	H	α MeI 0.36 2.24 4.1
CPK 197	"	2-Br	H	CH <sub>3</sub>	H	β MeI 0.24 0.50 28
CPK 155	"	3-Br	H	CH <sub>3</sub>	H	α MeI 2.0 0.8 49
CPK 184	"	3-Br	H	CH <sub>3</sub>	H	β MeI 0.31 0.11 7.7
CPK 165	"	4-Br	H	CH <sub>3</sub>	H	α MeI 0.95 0.62 61
CPK 171	"	4-Br	H	CH <sub>3</sub>	H	β MeI 0.21 0.14 29



TABLE 1 cont.

Compound	Inhibition of the uptake in vitro IC <sub>50</sub> (μM)					Inhibition of the uptake in vivo i.p. ED <sub>50</sub> (μmole/kg)				
	NA	5-HT	NA	5-HT	NA	5-HT	NA	5-HT	NA	5-HT
Compounds of the invention										
$  \begin{array}{c}  \text{Ar} \\    \\  \text{---CH-CH}_2\text{---CH-NHR}^1 \\    \\  \text{Ar}  \end{array}  $	Y	X	R	R <sup>1</sup>	isomer	salt <sup>1</sup>				
$  \begin{array}{c}  \text{Ar} \\    \\  \text{---CH-CH}_2\text{---CH-NHR}^1 \\    \\  \text{Ar}  \end{array}  $	4-Br	H	CH <sub>3</sub>	CH <sub>3</sub>	α+β	Ux	8.00	0.71	>98	98
PK 187										
PK 195	"	H	CH <sub>3</sub>	H	α	Mal	4.0	2.7	>98	98
PK 199	"	H	CH <sub>3</sub>	H	β	Mal	2.25	0.76	>98	>98
PK 160	"	H	CH <sub>3</sub>	H	α	Mal	0.51	0.30	78	54
PK 198	"	H	CH <sub>3</sub>	H	β	Mal	0.83	0.16	52	33
LA 615	"	4-F	CH <sub>3</sub>	H	-		1.9	0.30	57	29
LA 619	"	3-CH <sub>3</sub>	CH <sub>3</sub>	H	-		0.8	0.7	48	116
LA 614	"	4-CH <sub>3</sub>	CH <sub>3</sub>	H	-		0.80	0.15	100	25



Claim 1-3, 19

Not 45, 6, 2024

0000322


-31-

0000322

TABLE I cont.

Compound	Inhibition of the uptake in vitro IC <sub>50</sub> (μM)	Inhibition of the uptake in vivo i.p. ED <sub>50</sub> (μmole/kg)
	NA	5-HT
	5-HT	NA

## Compounds of the invention

Ar	Y	X	R	R <sup>1</sup>	isomer* salt**			
	4-Br	4-Br	CH <sub>3</sub>	H	-	HCl	4.8	2.5
	4-OCH <sub>3</sub>	4-OCH <sub>3</sub>	CH <sub>3</sub>	H	-	Fum	1.7	0.4
	4-F	4-F	CH <sub>3</sub>	H	-	Fum	1.5	0.3
	3-CH <sub>3</sub>	3-CH <sub>3</sub>	CH <sub>3</sub>	H	-	HCl	2.4	1.3
	4-Br	H	CH <sub>3</sub>	H	Z	Mal	2.0	3.0
	4-Br	H	CH <sub>3</sub>	H	E	Ox	2.6	0.6
	3-Br	H	CH <sub>3</sub>	H	Z	Mal	1.1	0.6
	3-Br	H	CH <sub>3</sub>	H	E	HCl	1.3	0.7
	3-pyridyl	4-Br	CH <sub>3</sub>	H	Z	Ox	1.6	0.2

—

TABLE I cont.

Compound	Inhibition of the uptake in vitro IC <sub>50</sub> (μM)		Inhibition of the uptake in vivo i.p. ED <sub>50</sub> (μmole/kg)	
	NA	5-HT	NA	5-HT
<u>Prior art compounds</u>				
3,3-diphenyl-1-methylpropylamine fumarate	0.50	0.50	50	75
1-methyl-3-(4-bromophenyl)-3-(3-pyridyl)-allylamine	0.8	2.5	25	102
-"-	1.5	0.10	>102	19
Chloroimipramine	0.9	0.09	150	20
Bromfeniramine	3.8	0.2	50	10

-33-

0000322

# Geometrical isomerism

## HCl = hydrochloride

Mal = maleate

Ox = oxalate

Fum = fumarate

-34-

0000322


The pharmacological tests show that the compounds are able to inhibit the uptake of noradrenaline and 5-hydroxytryptamine. A pronounced non-selective activity is shown for the compounds having codes CPK 170, CPK 171, CPK 180 and 5 CPK 184. A pronounced selective activity on uptake of noradrenaline is seen in compounds CPK 185, CPK 215 and CPK 217 while a pronounced selective activity on uptake of 5-hydroxytryptamine is seen in compounds FLA 611, FLA 615, CPK 198 and CPK 204.

10


A strong non-selective activity is considered to be especially advantageous, as compounds having such activity may be employed in the treatment of depressions in which the neurotransmitter deficiency is unknown as well as in those cases wherein 15 it is established that the deficiency pertains to both noradrenaline and 5-hydroxytryptamine.

The invention is summarized in the following clauses:

20 A. A compound of the general formula



25 wherein the dotted line represents an optional double bond and Ar represents the group



30


wherein Y is bound in the 2-, 3- or 4-position and represents a lower alkyl or lower alkoxy group, a halogen, a trifluoromethyl group or an amino, mono- or di-lower alkylamino group, or Ar represents a pyridyl group bound in the 2-, 3- or 4- 35 position, X represents hydrogen, a lower alkyl or lower alkoxy group, a halogen, a trifluoromethyl group or an amino, mono- or di-lower alkylamino group, R represents a lower alkyl group, and R<sup>1</sup> represents hydrogen or a lower alkyl group, whereby "lower" indicates a group having up to 3 carbon atoms; or a

-35-

0000322


therapeutically acceptable acid addition salt thereof, in anhydrous form or in any degree of hydration possible, or a bioprecursor thereof.

5 B. A compound of the general formula



10 wherein Ar, X, R, and R<sup>1</sup> have the meaning defined in clause A, or a salt or bioprecursor thereof as defined in clause A.

C. A compound of the general formula



wherein Ar, X, R and R<sup>1</sup> have the meaning defined in clause A, or a salt or bioprecursor thereof as defined in clause A.

20

D. A compound according to clause B in the form of a pure diastereomer.

E. A compound according to clause C in the form of a pure  
25 geometrical isomer.

F. A compound according to clause A in the form of a pure optical enantiomer.

30 G. A compound according to clause A wherein Ar represents the group



35

-35-

0000322

H. A compound according to clause G wherein X represents hydrogen.

J. A compound according to clause H wherein Y represents F,  
5 Br, or CH<sub>3</sub>O-.

K. A compound selected from the group consisting of:


- (β)-3-(4-fluorophenyl)-1-methyl-3-phenylpropylamine,  
10 (β)-3-(4-bromophenyl)-1-methyl-3-phenylpropylamine,  
(α)-3-(4-methoxyphenyl)-1-methyl-3-phenylpropylamine,  
(β)-3-(3-bromophenyl)-1-methyl-3-phenylpropylamine,  
(α)-3-(2-bromophenyl)-1-methyl-3-phenylpropylamine,  
(E)-3-amino-1-(3-bromophenyl)-1-phenylbutene,  
15 (Z)-3-amino-1-(3-bromophenyl)-1-phenylbutene,  
3-amino-1,1-di-(4-methoxyphenyl)-1-butene,  
3,3-di-(4-fluorophenyl)-1-methylpropylamine,  
(β)-3-(4-methoxyphenyl)-1-methyl-3-phenylpropylamine, and  
(E)-3-amino-1-(4-bromophenyl)-1-phenylbutene,  
20 or a salt or bioprecursor thereof as defined in clause A.

L. A compound according to one or more of the preceding clauses wherein R is a methyl group.

25 M. A compound according to one or more of the preceding clauses wherein R<sup>1</sup> is hydrogen.

N. A process for preparing a compound of the general formula


30



I

wherein the dotted line represents an optional double bond and Ar represents the group

35




-37-

0000322


wherein Y is bound in the 2-, 3-, or 4-position and represents a lower alkyl or lower alkoxy group, a halogen, a trifluoromethyl group or an amino, mono- or di-lower alkylamino group, or Ar represents a pyridyl group bound in the 2-, 3-, or 4-  
 5 position, X represents hydrogen, a lower alkyl or lower alkoxy group, a halogen, a trifluoromethyl group or an amino, mono- or di-lower alkylamino group, R represents a lower alkyl group, and R<sup>1</sup> represents hydrogen or a lower alkyl group, whereby "lower" indicates a group having up to 3 carbon atoms, character  
 10 ized in

a) reducing a compound of the formula




wherein Ar, X and R have the meaning defined in clause A and R<sup>1</sup> is a hydrogen atom or an alkyl, an acyl or an alkylsulfonyl group having 1-3 carbon atoms, to the obtention of a compound  
 20 of formula Ia, in which R<sup>1</sup> is hydrogen, or

b) preparing a reactive ester of an alcohol of the formula



wherein Ar, X, and R have the meaning defined in clause A, and reacting the ester obtained with an amine of the formula NH<sub>2</sub>R<sup>1</sup>, wherein R<sup>1</sup> has the meaning defined in clause A, or  
 30

c) reducing a compound of the formula




wherein Ar, X, R, and R<sup>1</sup> have the meaning defined in clause A or

-38-


0000322

d) reacting a ketone of the formula




wherein Ar, X, and R have the meaning defined in clause A, with ammoniumformate or methylammoniumformate according to Leuckart-Wallach,

10 e) dehydrating a carbinol of the formula




to a compound of formula Ib.

f) reacting a compound of the formula



wherein Ar, X, and R have the meaning defined above and Z is a leaving group with an amine of the formula  $\text{NH}_2\text{R}^1$  or a derivative thereof to the formation of a primary or secondary amine of formula Ib above, which formation involves conversion of an amine derivative when a derivative of an amine of the formula  $\text{NH}_2\text{R}^1$  is employed, or

30 g) oxidizing the benzylic carbon atom of a compound of the formula




wherein Ar, X, and R have the meaning defined above and R'' and R''' are protective groups for the amino function, followed by elimination of the group formed by oxidation at the benzyl



-39-


0000322

carbon atom, giving a compound of the formula



which is transformed into the compound of formula Ib by splitting; and if desired converting a primary amine obtained according to any process alternatives a) to g) above into the corresponding secondary amine by methylation, and if desired converting the primary or secondary amine obtained into a therapeutically acceptable acid addition salt thereof in anhydrous form or any degree of hydration possible.


15 P. A compound useful as an intermediate in preparation of the compound defined in clause A, characterized by the formula



wherein Ar, X, and R have the meaning defined in clause A and R' is a hydrogen atom or an alkyl, an acyl or an alkylsulfonyl group having 1-3 carbon atoms.


25

Q. A compound useful as an intermediate in preparation of the compound defined in clause A, characterized by the formula



wherein Ar, X, and R have the meaning defined in clause A.

35 S. A compound useful as an intermediate in preparation of the compound defined in clause A, characterized by the formula




-40-

0000322

wherein Ar, X, and R have the meaning defined in clause A.

T. A compound useful as an intermediate in preparation of the compound defined in claim 1, characterized by the formula

5




VII

10 wherein Ar, X, R, and R<sup>1</sup> have the meaning defined in clause A,

U. A compound useful as an intermediate in preparation of the compound defined in clause A, characterized by the formula

15




VIII

wherein Ar, X, R, and R<sup>1</sup> have the meaning defined in clause A  
20 and Z represents F, Cl, Br, I, OSO<sub>2</sub>R; wherein R<sup>1</sup> is an alkyl, aralkyl, or aryl group.

V. A compound useful as an intermediate in preparation of the compound defined in clause A, characterized by the formula

25




IX

30 wherein Ar, X, and R have the meaning defined in clause A, and R'' and R''' are protective groups for the amino function.

AA. A compound useful as an intermediate in preparation of the compound defined in clause A, characterized by the formula

35



X


-41-

0000322

wherein Ar, X, and R have the meaning defined in clause A.

AB. A compound useful as an intermediate in preparation of  
the compound defined in clause A, characterized by the formula

5



10 wherein Ar, X, and R have the meaning defined in clause A.

AC. A pharmaceutical preparation which comprises as active  
ingredient a therapeutically effective amount of a compound  
defined in clause A or a pharmaceutically acceptable salt  
15 thereof, in association with a pharmaceutically acceptable  
carrier.

AD. A method for the treatment of depressions, characterized  
in administration to a host suffering from such ailment a  
20 therapeutically acceptable amount of a compound defined in  
clause A or a pharmaceutically acceptable salt thereof.

AE. A method for the treatment of anxiety, characterized in  
administration to a host suffering from such ailment a thera-  
25apeutically acceptable amount of a compound defined in clause A  
or a pharmaceutically acceptable salt thereof.

AF. A compound, a process for preparing a compound, a compound  
useful as an intermediate, a pharmaceutical preparation and a  
30 method of treatment as defined in any of clauses A to AF and  
substantially as described.


BAD ORIGINAL

-1-


0000322

WHAT WE CLAIM IS:

1. A compound of the general formula



wherein the dotted line represents an optional double bond and  
10 Ar represents the group



15 wherein Y is bound in the 2-, 3-, or 4-position and represents  
a lower alkyl or lower alkoxy group, a halogen, a trifluoro-  
methyl group or an amino, mono- or di-lower alkylamino group,  
or Ar represents a pyridyl group bound in the 2-, 3-, 4-  
position, X represents hydrogen, a lower alkyl or lower al-  
20 koxy group, a halogen, a trifluoromethyl group or an amino,  
mono- or di-lower alkylamino group, R represents a lower alkyl  
group, and R<sup>1</sup> represents hydrogen or a lower alkyl group,  
whereby "lower" indicates a group having up to 3 carbon atoms;  
or a therapeutically acceptable acid addition salt thereof, in  
25 anhydrous form or in any degree of hydration possible, or a  
bioprecursor thereof.

2. A compound according to claim 1 in the form of

- 30 (a) a pure diastereomer,  
(b) a pure geometrical isomer, or  
(c) a pure optical enantiomer.

35

3. A compound according to claim 1 wherein Ar represents the  
group




-3-

0000322

di-lower alkylamino group, R represents a lower alkyl group, and R<sup>1</sup> represents hydrogen or a lower alkyl group, whereby "lower" indicates a group having up to 3 carbon atoms, characterized in

5


a) reducing a compound of the formula



wherein Ar, X, and R have the meaning defined in claim 1 and R' is a hydrogen atom or an alkyl, an acyl or an alkylsulfonyl group having 1-3 carbon atoms; to the obtention of a compound of formula Ia, in which R<sup>1</sup> is hydrogen, or

15


b) preparing a reactive ester of an alcohol of the formula



wherein Ar, X, and R have the meaning defined in claim 1, and reacting the ester obtained with an amine of the formula NH<sub>2</sub>R<sup>1</sup>, wherein R<sup>1</sup> has the meaning defined in claim 1, or


25

c) reducing a compound of the formula



wherein Ar, X, R, and R<sup>1</sup> have the meaning defined in claim 1 or


35 d) reacting a ketone of the formula



-2-

0000322

4. A compound according to claim 3 wherein X represents hydrogen.
5. A compound according to claim 4 wherein Y represents F, Br, or  $\text{CH}_3\text{O}-$ .
6. A compound selected from the group consisting of:
- ( $\beta$ )-3-(4-fluorophenyl)-1-methyl-3-phenylpropylamine,
  - 10 ( $\beta$ )-3-(4-bromophenyl)-1-methyl-3-phenylpropylamine,
  - ( $\alpha$ )-3-(4-methoxyphenyl)-1-methyl-3-phenylpropylamine,
  - ( $\beta$ )-3-(3-bromophenyl)-1-methyl-3-phenylpropylamine,
  - ( $\alpha$ )-3-(2-bromophenyl)-1-methyl-3-phenylpropylamine,
  - (E)-3-amino-1-(3-bromophenyl)-1-phenylbutene,
  - 15 (Z)-3-amino-1-(3-bromophenyl)-1-phenylbutene,
  - 3-amino-1,1-di-(4-methoxyphenyl)-1-butene,
  - 3,3-di-(4-fluorophenyl)-1-methylpropylamine,
  - ( $\beta$ )-3-(4-methoxyphenyl)-1-methyl-3-phenylpropylamine, and
  - (E)-3-amino-1-(4-bromophenyl)-1-phenylbutene
- 20 or a salt or bioprecursor thereof as defined in claim 1.
7. A process for preparing a compound of the general formula



wherein the dotted line represents an optional double bond and Ar represents the group



wherein Y is bound in the 2-, 3-, or 4-position and represents


35 a lower alkyl or lower alkoxy group, a halogen, a trifluoromethyl group or an amino, mono- or di-lower alkylamino group, or Ar represents a pyridyl group bound in the 2-, 3-, or 4-position, X represents hydrogen, a lower alkyl or lower alkoxy group, a halogen, a trifluoromethyl group or an amino, mono- or

5-

0000322

which is transformed into the compound of formula Ib by splitting; and if desired converting a primary amine obtained according to any of process alternatives a) to g) above into the corresponding secondary amine by methylation, and if desired, 5 converting the primary or secondary amine obtained into a therapeutically acceptable acid addition salt thereof in anhydrous form or any degree of hydration possible.


8. Compounds useful as intermediates in preparation of the 10 compound defined in claim 1, characterized by the formula



15

wherein Ar, X, and R have the meaning defined in claim 1 and R' is a hydrogen atom or an alkyl, an acyl or an alkylsulfonyl group having 1-3 carbon atoms;


20



25

wherein Ar, X, and R have the meaning defined in claim 1;

30




wherein Ar, X, and R have the meaning defined in claim 1;

-4-

0000322

wherein Ar, X, and R have the meaning defined in claim 1, with ammoniumformate or methylammoniumformate according to Leuckart-Wallach,

5 e) dehydrating a carbinol of the formula




VII

10

to a compound of formula Ib.

f) reacting a compound of the formula

15




VIII

wherein Ar, X, and R have the meaning defined above and Z is a leaving group with an amine of the formula  $\text{NH}_2\text{R}^1$  or a derivative thereof to the formation of a primary or secondary amine of formula Ib above, which formation involves conversion of an amine derivative when a derivative of an amine of the formula  $\text{NH}_2\text{R}^1$  is employed, or

25


g) oxidizing the benzylic carbon atom of a compound of the formula



IX

wherein Ar, X, and R have the meaning defined above and R'' and R''' are protective groups for the amino function, followed by elimination of the group formed by oxidation at the benzylic carbon atom, giving a compound of the formula

40



X

BAD ORIGINAL



-7-


00003225

wherein Ar, C, and R have the meaning defined in claim 1.


9. A pharmaceutical preparation which comprises as active ingredient a therapeutically effective amount of a compound defined in claim 1 or a pharmaceutically acceptable salt thereof in association with a pharmaceutically acceptable carrier.

-6-


0000322




5 wherein Ar, X, R and R<sup>1</sup> have the meaning defined in claim 1;




wherein Ar, X, R, and R<sup>1</sup> have the meaning defined in claim 1  
 and Z represents F, Cl, Br, I, OSO<sub>2</sub>R', wherein R' is an alkyl,  
 15 aralkyl, or aryl group;



wherein Ar, X, and R have the meaning defined in claim 1 and R''  
 and R''' are protective groups for the amino function;  
 25



wherein Ar, X, and R have the meaning defined in claim 1;



BAD ORIGINAL



European Patent  
Office

# 0000322 EUROPEAN SEARCH REPORT

Application number  
EP 78 85 000

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	CHEMICAL ABSTRACTS, vol. 82, (1975) nr. 170039c As intermediate : 3,3-di-(4-fluorophenyl)-1-methyl-propylamine	1-3,5-8	C 07 C 87/28 C 07 C 87/29 C 07 C 93/14 C 07 C 49/16 C 07 C 131/00 C 07 C 49/06 C 07 C 91/18 C 07 D 209/48 C 07 D 213/38 A 61 K 31/135 A 61 K 31/44 // C 07 C 125/06 C 07 C 25/26
A	GB - A - 624 117 (THE WELLCOME FOUNDATION) * Pages 5-6; claims *	1,7	TECHNICAL FIELDS SEARCHED (Int. Cl.)  C 07 C 87/28 C 07 C 87/29 C 07 C 93/14 C 07 D 213/38   CATEGORY OF CITED DOCUMENTS X: particularly relevant A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: conflicting application D: document cited in the application L: citation for other reasons  G: member of the same patent family, corresponding document
A	FR - A - 1 021 301 (THE LUNDBECK CORPORATION) * Pages 4-5; abstract *	1,7	
A	GB - A - 955 516 (N.V. KONINKLIJKE PHARMACEUTISCHE FABRIEKEN v/h BROCADES - STHEEMAN & PHARMACIA) * Page 11; claims *	1,7	
A	FR - A - 1 160 995 (FARBWERKE HOECHST A.G.) * Page 5, abstract *	1,7	
A	FR - A - 1 226 401 (M. ERLBACH AND A. SIEGLITZ) * Page 5, abstract *	1,7	
<input checked="" type="checkbox"/> The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of the search 10-10-1978	Examiner PAUWELS